

AD-A125 245

LINEAR PARABOLIC EQUATIONS WITH A SINGULAR LOWER ORDER  
COEFFICIENT(U) WISCONSIN UNIV-MADISON MATHEMATICS  
RESEARCH CENTER K HOELLIG NOV 82 MRC-TSR-2453


1/1


UNCLASSIFIED

DAG29-80-C-0041

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD A125245

MRC Technical Summary Report #2453


LINEAR PARABOLIC EQUATIONS WITH A  
SINGULAR LOWER ORDER COEFFICIENT

Klaus Höllich

Mathematics Research Center  
University of Wisconsin-Madison  
610 Walnut Street  
Madison, Wisconsin 53706

November 1982

(Received September 28, 1982)



Approved for public release  
Distribution unlimited

Sponsored by

U. S. Army Research Office  
P. O. Box 12211  
Research Triangle Park  
North Carolina 27709

National Science Foundation  
Washington, DC 20550

83 02 023 120

DTIC FILE COPY

UNIVERSITY OF WISCONSIN - MADISON  
MATHEMATICS RESEARCH CENTER

LINEAR PARABOLIC EQUATIONS WITH A SINGULAR  
LOWER ORDER COEFFICIENT

Klaus Hölzig

Technical Summary Report #2453

November 1982

ABSTRACT

Let  $a, b$  belong to the Hölder class  $H^{\alpha, \alpha/2}([0,1] \times [0,T])$  with  $\alpha \in (0,1)$  and  $a > a_0 > 0$ . It is shown that for the solution of the problem

$$u_t - au_{xx} - t^{-1/2}bu_x = t^{-1/2}f, \quad (x,t) \in [0,1] \times [0,T],$$

$$u(\cdot, 0) = \phi$$

$$u(v, \cdot) = \psi_v, \quad v = 0, 1,$$

the estimate

$$\|u, u_x, t^{1/2}u_{xx}\|_{\alpha, \alpha/2} <$$

$$c\{\|f\|_{\alpha, \alpha/2} + \|\phi, \phi'\|_{\alpha} + \sum_{v=0}^1 \|\psi_v, t^{1/2}\psi'_v\|_{\alpha/2}\}$$

holds if the data  $f, \phi, \psi_v$  satisfy the appropriate compatibility conditions.

Here  $\|\cdot\|_{\alpha, \alpha/2}, \|\cdot\|_{\alpha}, \|\cdot\|_{\alpha/2}$  denote the norms of the Hölder classes

$H^{\alpha, \alpha/2}([0,1] \times [0,T]), H^{\alpha}([0,1])$  and  $H^{\alpha/2}([0,T])$  respectively and the

constant  $c$  depends on  $\alpha, a_0, \|a, b\|_{\alpha, \alpha/2}$  and  $T$ . The result extends to quasilinear problems with  $a, b$  and  $f$  depending on  $x, t$  and  $u$ .

AMS (MOS) Subject Classification: 35K20

Key Words: parabolic equations, linear, singular, regularity

Work Unit Number 1 - Applied Analysis

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041. This material is based upon work supported by the National Science Foundation under Grant No. MCS-7927062, Mod. 2.

### SIGNIFICANCE AND EXPLANATION

This report was motivated by the study of free boundary value problems related to the Stefan problem. The precise description of the smoothness of the free boundary requires sharp regularity results for linear parabolic equations with singular coefficients.

These results are of independent interest. They can be applied to parabolic equations on domains with curved boundaries that touch the x-axis. As a particular example consider the heat equation ;

$$u_t - u_{xx} = t^{-1/2} f \quad \text{on } \Omega$$

$$u = 0 \quad \text{on } \partial\Omega$$

on the domain  $\Omega := \{(x,t) : t > 0, x > -t^{1/2}\}$ . Our results imply that  $u$  and  $u_x$  are Hölder continuous up to the boundary if  $f$  is and  $f(0,0) = 0$ . This example arises in the *convexity* of the nonlinear parabolic problem studied in MRC Technical Summary Report #2354.

Accession For

NEWS ORAL ☒

FILED ☐

U. S. ☐

1. ☐

2. ☐

3. ☐

4. ☐

5. ☐

6. ☐

7. ☐

8. ☐

9. ☐

10. ☐

11. ☐

12. ☐

13. ☐

14. ☐

15. ☐

16. ☐

17. ☐

18. ☐

19. ☐

20. ☐

21. ☐

22. ☐

23. ☐

24. ☐

25. ☐

26. ☐

27. ☐

28. ☐

29. ☐

30. ☐

31. ☐

32. ☐

33. ☐

34. ☐

35. ☐

36. ☐

37. ☐

38. ☐

39. ☐

40. ☐

41. ☐

42. ☐

43. ☐

44. ☐

45. ☐

46. ☐

47. ☐

48. ☐

49. ☐

50. ☐

51. ☐

52. ☐

53. ☐

54. ☐

55. ☐

56. ☐

57. ☐

58. ☐

59. ☐

60. ☐

61. ☐

62. ☐

63. ☐

64. ☐

65. ☐

66. ☐

67. ☐

68. ☐

69. ☐

70. ☐

71. ☐

72. ☐

73. ☐

74. ☐

75. ☐

76. ☐

77. ☐

78. ☐

79. ☐

80. ☐

81. ☐

82. ☐

83. ☐

84. ☐

85. ☐

86. ☐

87. ☐

88. ☐

89. ☐

90. ☐

91. ☐

92. ☐

93. ☐

94. ☐

95. ☐

96. ☐

97. ☐

98. ☐

99. ☐

100. ☐

101. ☐

102. ☐

103. ☐

104. ☐

105. ☐

106. ☐

107. ☐

108. ☐

109. ☐

110. ☐

111. ☐

112. ☐

113. ☐

114. ☐

115. ☐

116. ☐

117. ☐

118. ☐

119. ☐

120. ☐

121. ☐

122. ☐

123. ☐

124. ☐

125. ☐

126. ☐

127. ☐

128. ☐

129. ☐

130. ☐

131. ☐

132. ☐

133. ☐

134. ☐

135. ☐

136. ☐

137. ☐

138. ☐

139. ☐

140. ☐

141. ☐

142. ☐

143. ☐

144. ☐

145. ☐

146. ☐

147. ☐

148. ☐

149. ☐

150. ☐

151. ☐

152. ☐

153. ☐

154. ☐

155. ☐

156. ☐

157. ☐

158. ☐

159. ☐

160. ☐

161. ☐

162. ☐

163. ☐

164. ☐

165. ☐

166. ☐

167. ☐

168. ☐

169. ☐

170. ☐

171. ☐

172. ☐

173. ☐

174. ☐

175. ☐

176. ☐

177. ☐

178. ☐

179. ☐

180. ☐

181. ☐

182. ☐

183. ☐

184. ☐

185. ☐

186. ☐

187. ☐

188. ☐

189. ☐

190. ☐

191. ☐

192. ☐

193. ☐

194. ☐

195. ☐

196. ☐

197. ☐

198. ☐

199. ☐

200. ☐

201. ☐

202. ☐

203. ☐

204. ☐

205. ☐

206. ☐

207. ☐

208. ☐

209. ☐

210. ☐

211. ☐

212. ☐

213. ☐

214. ☐

215. ☐

216. ☐

217. ☐

218. ☐

219. ☐

220. ☐

221. ☐

222. ☐

223. ☐

224. ☐

225. ☐

226. ☐

227. ☐

228. ☐

229. ☐

230. ☐

231. ☐

232. ☐

233. ☐

234. ☐

235. ☐

236. ☐

237. ☐

238. ☐

239. ☐

240. ☐

241. ☐

242. ☐

243. ☐

244. ☐

245. ☐

246. ☐

247. ☐

248. ☐

249. ☐

250. ☐

251. ☐

252. ☐

253. ☐

254. ☐

255. ☐

256. ☐

257. ☐

258. ☐

259. ☐

260. ☐

261. ☐

262. ☐

263. ☐

264. ☐

265. ☐

266. ☐

267. ☐

268. ☐

269. ☐

270. ☐

271. ☐

272. ☐

273. ☐

274. ☐

275. ☐

276. ☐

277. ☐

278. ☐

279. ☐

280. ☐

281. ☐

282. ☐

283. ☐

284. ☐

285. ☐

286. ☐

287. ☐

288. ☐

289. ☐

290. ☐

291. ☐

292. ☐

293. ☐

294. ☐

295. ☐

296. ☐

297.

**The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.**

## Table of Contents

1.	<u>Statement of the result</u>	1
2.	<u>The constant coefficient problem</u>	4
2.1	Auxiliary lemmas	4
2.2	The Cauchy problem	6
2.3	The boundary value problem	10
3.	<u>Extension to variable coefficients</u>	22
3.1	Reduction of the problem	22
3.2	Auxiliary lemmas	24
3.3	Proof of the main theorem	25

# LINEAR PARABOLIC EQUATIONS WITH A SINGULAR LOWER ORDER COEFFICIENT

Klaus Hölzig

## 1. Statement of the result

We consider the linear parabolic initial boundary value problem

$$(1.1) \quad \begin{cases} u_t(x,t) - a(x,t)u_{xx}(x,t) - t^{-1/2}b(x,t)u_x(x,t) = t^{-1/2}f(x,t), \\ (x,t) \in \Omega_T := [0,1] \times [0,T], \\ u(x,0) = \phi(x), \quad x \in [0,1], \\ u(v,t) = \psi_v(t), \quad t \in [0,T], \quad v = 0,1. \end{cases}$$

Problems of this type, with a singular coefficient of  $u_x$ , may arise when transforming parabolic equations from a domain with curved boundaries to the standard domain  $\Omega_T$ . Consider, e.g., the heat equation,

$$(1.2) \quad \begin{cases} v_t - v_{yy} = t^{-1/2}g & \text{on } \Omega \\ v = 0 & \text{on } \partial\Omega \end{cases}$$

on the domain  $\Omega = \{(y,t) : 0 < y < 1 + t^{1/2}, t \in [0,T]\}$ . By the change of variables  $x = y/(1 + t^{1/2})$ , (1.2) is equivalent to (1.1) with  $u(x,t) = v(y,t)$ ,  $f(x,t) = g(y,t)$ ,  $\phi = \psi_v = 0$ ,  $a = (1 + t^{1/2})^{-2}$  and  $b(x,t) = \frac{1}{2}x/(1 + t^{1/2})$ . More generally, any parabolic equation on a domain with smooth vertical boundaries that touch the  $x$ -axis, but have nonzero curvature as  $t \rightarrow 0$ , leads, after a change of variables, to a problem of the form (1.1).

We shall show that, under the assumptions on the coefficients and the data specified in (1.3) - (1.5) below, the solution of problem (1.1) and its partial derivative with respect to  $x$  are Hölder continuous up to the

boundary. This is, in general, no longer true for a singularity of the form  $t^{-1/2-\varepsilon}$ ,  $\varepsilon > 0$ , in the coefficient of  $u_x$ .

We use the Hölder norms

$$|\chi|_{\alpha, I} := \sup_{z, z' \in I} |\chi(z) - \chi(z')| / |z - z'|^\alpha$$

$$\|\chi\|_{\alpha, I} := \|\chi\|_{\infty, I} + |\chi|_{\alpha, I}$$

$$|w|_{\alpha, x, \Omega} := \sup_{(x, t), (x', t) \in \Omega} |w(x, t) - w(x', t)| / |x - x'|^\alpha$$

$$|w|_{\beta, t, \Omega} := \sup_{(x, t), (x, t') \in \Omega} |w(x, t) - w(x, t')| / |t - t'|^\beta$$

$$|w|_{\alpha, \beta, \Omega} := |w|_{\alpha, x, \Omega} + |w|_{\beta, t, \Omega}$$

$$\|w\|_{\alpha, \beta, \Omega} := \|w\|_{\infty, \Omega} + |w|_{\alpha, \beta, \Omega}$$

where  $\alpha, \beta \in (0, 1)$  and denote by  $H^\alpha(I)$  and  $H^{\alpha, \beta}(\Omega)$  the Hölder classes corresponding to the norms  $\|\cdot\|_{\alpha, I}$  and  $\|\cdot\|_{\alpha, \beta, \Omega}$ . The subscripts  $I$  and  $\Omega$  are omitted if the domains are clear from the context. We also need the subspaces  $\overset{\circ}{H}^\alpha([0, T]) := \{\chi \in H^\alpha([0, T]) : \chi(0) = 0\}$  and  $\overset{\circ}{H}^{\alpha, \beta}(I \times [0, T]) := \{w \in H^{\alpha, \beta}(I \times [0, T]) : w(x, 0) = 0, x \in I\}$ . For simplicity of notation we write  $t^\gamma \chi$  and  $t^\gamma w$  for the functions  $t \mapsto t^\gamma \chi(t)$  and  $(x, t) \mapsto t^\gamma w(x, t)$ .

We assume that the coefficients  $a, b$  and the data  $f, \phi, \psi_\nu$  for the problem (1.1) satisfy

$$(1.3) \quad \begin{cases} a, b \in H^{\alpha, \alpha/2} \\ a > a_0 > 0 \end{cases}$$

$$(1.4) \quad \begin{cases} f \in H^{\alpha, \alpha/2} \\ \phi, \phi' \in H^\alpha \\ \psi_\nu, t^{1/2} \psi'_\nu \in H^{\alpha/2} \end{cases}$$

for some  $\alpha \in (0,1)$  and the compatibility conditions

$$(1.5) \quad \begin{cases} \phi(v) = \psi_v(0) , \\ \lim_{t \rightarrow 0} t^{1/2} \psi'_v(t) = f(v,0) + b(v,0)\phi'(v), \quad v = 0,1 . \end{cases}$$

Theorem 1.1. Under the assumptions (1.3) - (1.5) the solution of the problem (1.1) satisfies

$$(1.6) \quad \|u, u_x, t^{1/2} u_{xx}\|_{\alpha, \alpha/2, \Omega_T} \leq c \{ \|f\|_{\alpha, \alpha/2, \Omega_T} + \|\phi, \phi'\|_{\alpha, [0,1]} + \sum_{v=0}^1 \|\psi_v, t^{1/2} \psi'_v\|_{\alpha/2, [0,T]} \}$$

where the constant  $c$  depends on  $\alpha, a_0, \|a, b\|_{\alpha, \alpha/2, \Omega_T}$  and  $T$ . Moreover we have

$$(1.7) \quad \lim_{t \rightarrow 0} t^{1/2} u_{xx}(x,t) = 0, \quad x \in [0,1] .$$

The Theorem remains valid if we replace in problem (1.1)  $t^{-1/2}b(x,t)$  by  $t^{-\gamma}b(x,t)$  with  $\gamma < 1/2$ . In this case  $t^{-\gamma}b u_x$  can be regarded as a minor term and the result can be obtained from Theorem 1.1 by iteration. The Theorem also extends to quasilinear problems of the type (1.1) with  $a = a(x,t,u)$ ,  $b = b(x,t,u)$ ,  $f = f(x,t,u)$  if we assume, e.g., that  $a, b, f$  are  $C^1$ . We stated the result in the simplest setting to focus on its essential feature, the singularity in the coefficient of  $u_x$ .

The difficult part of the proof of Theorem 1.1 is to show (1.6) for the constant coefficient problem on the domain  $R_+ \times [0,T]$  (section 2.3). We then use a standard technique [L, pp. 295-341] to extend the result to the case of variable coefficients (section 3). To keep the report self contained, the proofs of estimates for the Cauchy problem (section 2.2) are included

although they are similar to the corresponding estimates for the heat equation.

## 2. The constant coefficient problem

### 2.1. Auxiliary lemmas

We list in this section some elementary inequalities and properties of Hölder norms. In the sequel  $c$  denotes generic constants which may depend on  $\alpha, \gamma, a_0, |a, b|_{\alpha, \alpha/2}$ .

(2.1) For  $z, j > 0$ ,

$$z^j \exp(-z) < c \exp(-cz) .$$

(2.2) For  $t > s > 0, z \in \mathbb{R}$ ,

$$-c + c|z|/(t-s)^{1/2} < |z + t^{1/2} - s^{1/2}|/(t-s)^{1/2} < c + c|z|/(t-s)^{1/2} .$$

(2.3) For  $t, z > 0, j > 1$ ,

$$\int_0^t s^{-j/2} (t-s)^{-1/2} \exp(-\frac{z}{s}) ds < cz^{(1-j)/2} .$$

Lemma 2.1. For  $\chi \in H^\gamma([0, T])$ ,  $\gamma \in (0, 1)$ ,

$$(2.4) \quad |\chi|_\gamma < c \sup_{j \in \mathbb{N}} |\chi|_{\gamma, [2^{-j}T, 2^{-j+1}T]} .$$

When estimating differences  $\chi(t) - \chi(t')$  it will be sometimes convenient to assume that  $|t - t'| < c \min(t, t')$ . In view of the Lemma this is no loss of generality when estimating  $|\chi|_\gamma$ .

Lemma 2.2. Let  $v(y, t) := u(x + ct^{1/2}, t)$ , then

$$(2.5) \quad |v|_{\alpha/2, t} < c |u|_{\alpha, \alpha/2} .$$

Lemma 2.3. For  $t^{1/2}\chi \in H^\gamma([0, T])$ ,  $\gamma \in (0, 1)$ , the norms  $\|t^{1/2}\chi\|_\gamma$ ,  $\|t^{1/2}\chi\|_\gamma$  and  $(\|t^{1/2-\gamma}\chi\|_\infty + \sup_{t \in (0, T)} t^{1/2} |\chi|_{\gamma, [t, T]})$  are equivalent.

An analogous version of this Lemma holds for  $\frac{\alpha}{H}, \alpha/2$ .

Proof. Assume that  $\|t^{1/2-\gamma} \chi\|_{\infty} + \sup t^{1/2} |\chi|_{\gamma, [t, T]} < 1$  and let  $0 < s < t < T$ . We have

$$\begin{aligned} & |t^{1/2} \chi(t) - s^{1/2} \chi(s)| < \\ & |(t^{1/2} - s^{1/2}) \chi(t)| + |s^{1/2} (\chi(t) - \chi(s))| < \\ & (t-s) t^{-1/2} t^{-1/2+\gamma} + (t-s)^{\gamma} < 2(t-s)^{\gamma} \end{aligned}$$

and therefore  $\|t^{1/2} \chi\|_{\gamma} < 2$ .

Now assume that  $\|\bar{\chi}\|_{\gamma} < 1$  with  $\bar{\chi} := t^{1/2} \chi$  and let  $0 < s/2 < t < s < T$ . We have

$$\begin{aligned} & |s^{-1/2} \bar{\chi}(s) - t^{-1/2} \bar{\chi}(t)| < \\ & |s^{-1/2} (\bar{\chi}(s) - \bar{\chi}(t))| + |(s^{-1/2} - t^{-1/2}) \bar{\chi}(t)| < \\ & s^{-1/2} (s-t)^{\gamma} + t^{-3/2} (s-t) t^{\gamma} < \\ & 2t^{-1/2} (s-t)^{\gamma} \end{aligned}$$

and therefore  $|\chi|_{\gamma, [t, T]} < ct^{-1/2}$ .

Lemma 2.4. For  $\gamma \in (0, 1/2)$  define

$$(2.6) \quad Q^{\gamma} := \{ \chi : \chi(0) = \lim_{t \rightarrow 0} t^{1/2} \chi'(t) = 0, \\ \|\chi\|_Q := \|t^{1/2} \chi'\|_{\gamma, [0, T]} < \infty \}.$$

Then we have, for  $\chi \in Q^{\gamma}$ ,

$$(2.7) \quad \|\chi\|_{\gamma+1/2}, \|t^{-1/2} \chi\|_{\gamma} < c \|\chi\|_Q.$$

## 2.2. The Cauchy problem

In this section we consider the problem

$$(2.8) \quad \begin{cases} u_t - au_{xx} - t^{-1/2}bu_x = t^{-1/2}f, & (x,t) \in \mathbb{R} \times [0,T] , \\ u(x,0) = \phi(x), & x \in \mathbb{R} , \end{cases}$$

where  $a > 0$  and  $b$  are constants.

Theorem 2.1. Let  $f \in H^{\alpha, \alpha/2}(\mathbb{R} \times [0,T])$ ,  $\phi, \phi' \in H^{\alpha}(\mathbb{R})$ . Then the solution of problem (2.8) satisfies

$$(2.9) \quad \|u, u_x, t^{1/2}u_{xx}\|_{\alpha, \alpha/2} < c\{\|f\|_{\alpha, \alpha/2} + \|\phi, \phi'\|_{\alpha}\}$$

where  $c$  depends on  $a$  and  $b$ . Moreover we have

$$(2.10) \quad \lim_{t \rightarrow 0} t^{1/2}u_{xx}(x,t) = 0, \quad x \in \mathbb{R} .$$

The change of variables  $y := x + 2bt^{1/2}$ ,  $v(y,t) := u(x,t)$ ,  $g(y,t) := f(x,t)$ , transforms the problem (2.8) into the heat equation

$$(2.11) \quad \begin{cases} v_t - av_{yy} = t^{-1/2}g, & (y,t) \in \mathbb{R} \times [0,T] , \\ v(y,0) = \phi(y), & y \in \mathbb{R} . \end{cases}$$

Therefore the analysis of the Cauchy problem is fairly simple. We merely have to take into account the singular behavior of  $t^{-1/2}g$  as  $t \rightarrow 0$ . However, for convenience of the reader, we give a complete proof of Theorem 2.1.

By Lemma 2.2 it suffices to prove the assertion (2.9) and (2.10) for the solution  $v$  of problem (2.11), i.e.

$$(2.12) \quad \|v, v_y, t^{1/2}v_{yy}\|_{\alpha, \alpha/2} < c\{\|g\|_{\alpha, \alpha/2} + \|\phi, \phi'\|_{\alpha}\}$$

$$(2.13) \quad \lim_{t \rightarrow 0} t^{1/2}v_{yy}(y,t) = 0, \quad y \in \mathbb{R} .$$

Denote by  $\Gamma(x,t) := (4\pi t)^{-1/2} \exp(-\frac{x^2}{4t})$  the fundamental solution of the heat equation. We have the estimate [L, p. 274]

$$(2.14) \quad |D_x^j D_t^k \Gamma(x,t)| \leq c t^{-j/2-k} \exp(-c \frac{x^2}{t}) .$$

Also note that

$$(2.15) \quad \int_{\mathbb{R}} D_x^j \Gamma(x,t) dx = \begin{cases} 1 & , j = 0 \\ 0 & , j > 0 \end{cases} .$$

$$(2.16) \quad \int_0^\infty \Gamma_x(x,t) dt = -\frac{1}{2} .$$

To prove (2.12) and (2.13) we assume that  $a = 1$  in problem (2.11) and consider two cases

(i) For  $\underline{q} = 0$  we have

$$\|v, v_y\|_{\alpha, \alpha/2} \leq c \|\phi, \phi'\|_{\alpha}$$

$$\|v_{yy}\|_{\alpha, \alpha/2, \mathbb{R}^+ \times [t, T]} \leq c \|v_{yy}(\cdot, t)\|_{\alpha} \leq c t^{-1/2} \|\phi'\|_{\alpha} .$$

By Lemma 2.3, this proves (2.12) once we show

$$\|t^{1/2-\alpha/2} v_{yy}\|_{\infty} \leq c \|\phi'\|_{\alpha}$$

which, in turn, yields (2.13). To this end we note that by (2.14) and (2.15),

$$|v_{yy}(y,t)| = \left| \int_{\mathbb{R}} \Gamma_y(y-z,t) (\phi'(z) - \phi'(y)) dz \right| \leq c t^{-1/2+\alpha/2} \|\phi'\|_{\alpha} .$$

(ii) Now assume that  $\underline{\phi} = 0$ .

Lemma 2.5.

$$(2.17) \quad \|v_y\|_{\alpha, y} \leq c \|g\|_{\alpha, y} .$$

Proof. Assume that  $\|g\|_{\alpha, y} < 1$  and let  $y-y' =: h > 0$ . Using (2.15) we have

$$\begin{aligned} v_y(y,t) - v_y(y',t) &= \\ \int_0^t \int_{|y-z| \leq 2h} \Gamma_y(y-z, t-s) s^{-1/2} (g(z,s) - g(y,s)) dz ds \end{aligned}$$

$$\begin{aligned}
& - \int_0^t \int_{|y-z| \leq 2h} \Gamma_y(y'-z, t-s) s^{-1/2} (g(z, s) - g(y', s)) dz ds \\
& + \int_0^t \int_{|y-z| \geq 2h} (\Gamma_y(y-z, t-s) - \Gamma_y(y'-z, t-s)) s^{-1/2} (g(z, s) - g(y, s)) dz ds \\
& + \int_0^t \int_{|y-z| \geq 2h} \Gamma_y(y'-z, t-s) s^{-1/2} (g(y', s) - g(y, s)) dz ds \\
& =: \sum_{\nu=1}^4 I_\nu.
\end{aligned}$$

Estimating these terms, using (2.14), we get

$$\begin{aligned}
|I_1| & \leq c \iint (t-s)^{-1} \exp(-c \frac{(y-z)^2}{t-s}) s^{-1/2} |y-z|^\alpha dz ds \\
& \leq c \int_{|y-z| \leq 2h} |y-z|^{-1+\alpha} dz \leq ch^\alpha.
\end{aligned}$$

For the second inequality we have used (2.3) with  $j = 2$ .  $|I_2|$  is estimated similarly.

$$|I_3| \leq h \iint |\Gamma_{yy}(\xi-z, t-s)| s^{-1/2} |z-y|^\alpha dz ds$$

with  $\xi = \xi(z, s) \in (y', y)$ . Since for  $|y-z| \geq 2h = 2(y-y')$ ,  $|y-z|/2 \leq |\xi-z| \leq 2|y-z|$ , we obtain, using (2.3) with  $j = 3$ ,

$$\begin{aligned}
|I_3| & \leq ch \iint (t-s)^{-3/2} \exp(-c \frac{(y-z)^2}{t-s}) s^{-1/2} |y-z|^\alpha dz ds \\
& \leq ch \int_{|z-y| \geq 2h} |y-z|^{-2+\alpha} dz \leq ch^\alpha. \\
|I_4| & \leq \int_0^t |\Gamma(y'-z, t-s)| \Big|_{z=y-2h}^{z=y+2h} s^{-1/2} h^\alpha ds \\
& \leq ch^\alpha \int_0^t (t-s)^{-1/2} s^{-1/2} ds \leq ch^\alpha.
\end{aligned}$$

Lemma 2.6.

$$(2.18) \quad |v_y|_{\alpha/2, t} \leq c |g|_{\alpha, y}.$$

Proof. Assume that  $|g|_{\alpha, y} < 1$  and let  $t-t' =: h \in (0, t/2)$ . Using (2.15) we have

$$\begin{aligned} v_y(y, t) - v_y(y, t') &= \\ &= \int_{t-2h}^t \int_R \Gamma_y(y-z, t-s) s^{-1/2} (g(z, s) - g(y, s)) dz ds \\ &- \int_{t-2h}^{t'} \int_R \Gamma_y(y-z, t'-s) s^{-1/2} (g(z, s) - g(y, s)) dz ds \\ &+ \int_0^{t-2h} \int_R (\Gamma_y(y-z, t-s) - \Gamma_y(y-z, t'-s)) s^{-1/2} (g(z, s) - g(y, s)) dz ds \\ &=: \sum_{\nu=1}^3 I_\nu. \end{aligned}$$

Estimating these terms, using (2.14), we get

$$\begin{aligned} |I_1| &< c \iint (t-s)^{-1} \exp\left(-c \frac{(y-z)^2}{t-s}\right) s^{-1/2} |y-z|^\alpha dz ds \\ &< c \int_{t-2h}^t (t-s)^{-1/2+\alpha/2} s^{-1/2} ds < ch^{\alpha/2}. \end{aligned}$$

$|I_2|$  is estimated similarly.

$$\begin{aligned} |I_3| &< h \iint |\Gamma_{yt}(y-z, t-s)| s^{-1/2} |y-z|^\alpha dz ds \\ &< ch \iint (t-s)^{-2} \exp\left(-c \frac{(y-z)^2}{t-s}\right) s^{-1/2} |y-z|^\alpha dz ds \\ &< ch \int_0^{t-2h} (t-s)^{-3/2+\alpha/2} s^{-1/2} ds < ch^{\alpha/2}. \end{aligned}$$

Lemma 2.7.

$$(2.19) \quad |t^{1/2-\alpha/2} v_{yy}|_\infty < c |g|_{\alpha, y}.$$

Proof. Assuming  $|g|_{\alpha, y} < 1$  we have, using (2.14) and (2.15),

$$\begin{aligned} |v_{yy}(y, t)| &= \left| \int_0^t \int_R \Gamma_{yy}(y-z, t-s) s^{-1/2} (g(z, s) - g(y, s)) dz ds \right| \\ &< c \iint (t-s)^{-3/2} \exp\left(-c \frac{(y-z)^2}{t-s}\right) s^{-1/2} |y-z|^\alpha dz ds \end{aligned}$$

$$< c \int_0^t (t-s)^{-1+\alpha/2} s^{-1/2} ds < ct^{-1/2+\alpha/2}.$$

This Lemma proves (2.13). In combination with Lemma 2.6 and the fact that  $v(\cdot, 0) = 0$  it also shows that

$$|v|_{1/2, t} < |t^{1/2} v_t|_{\infty} < c|g|_{\alpha, \alpha/2}$$

and it follows from Lemma 2.5 that

$$|v, v_y|_{\alpha, \alpha/2} < c|g|_{\alpha, \alpha/2}.$$

To complete the proof of assertion (2.12), we note that

$$\begin{aligned} & |v_{yy}|_{\alpha, \alpha/2, R \times [s, T]} < \\ & c(|v_{yy}(\cdot, s)|_{\alpha} + |t^{-1/2} g|_{\alpha, y, R \times [s, T]}) < \\ & c(s^{-1/2} |v_y(\cdot, s/2)|_{\alpha} + s^{-1/2} |g|_{\alpha, y, R \times [s/2, T]}) < \\ & c s^{-1/2} |g|_{\alpha, y}. \end{aligned}$$

In view of Lemmas 2.3 and 2.7 this yields

$$|t^{1/2} v_{yy}|_{\alpha, \alpha/2} < c|g|_{\alpha, y}.$$

### 2.3. The boundary value problem

In this section we consider the problem

$$(2.20) \quad \begin{cases} u_t - au_{xx} - t^{-1/2} bu_x = t^{-1/2} f, & (x, t) \in R_+ \times [0, T] \\ u(x, 0) = \phi(x), & x \in R_+ \\ u(0, t) = \psi(t), & t \in [0, T] \end{cases}$$

where  $a > 0$  and  $b$  are constants.

**Theorem 2.2.** Let  $f \in H^{\alpha, \alpha/2}(\mathbb{R}_+ \times [0, T])$ ,  $\phi, \phi' \in H^{\alpha}(\mathbb{R}_+)$ ,  $\psi, t^{1/2}\psi' \in H^{\alpha/2}([0, T])$  and assume that the compatibility conditions

$$(2.21) \quad \begin{cases} \phi(0) = \psi(0) \\ \lim_{t \rightarrow 0} t^{1/2} \psi(t) = f(0, 0) + b\phi'(0) \end{cases}$$

hold. Then the solution of problem (2.20) satisfies

$$(2.22) \quad \|u, u_x, t^{1/2}u_{xx}\|_{\alpha, \alpha/2} \leq c\{\|f\|_{\alpha, \alpha/2} + \|\phi, \phi'\|_{\alpha} + \|\psi, t^{1/2}\psi'\|_{\alpha/2}\}$$

where  $c$  depends on  $a$  and  $b$ . Moreover we have

$$(2.23) \quad \lim_{t \rightarrow 0} t^{1/2} u_{xx}(x, t) = 0, \quad x \in \mathbb{R}_+.$$

Denote by  $\bar{f}, \bar{\phi}$  smooth extensions of the function  $f, \phi$  to the domains  $\mathbb{R} \times [0, T]$  and  $\mathbb{R}$  respectively. Subtracting from  $u$  the solution of the Cauchy problem with right hand side  $t^{-1/2} \bar{f}$  and initial values  $\bar{\phi}$  and using Theorem 2.1, we see that we have to prove Theorem 2.2 only for  $f = \phi = 0$ , an assumption we make throughout this section. In this case, the compatibility conditions (2.21), together with the assumption  $\psi, t^{1/2}\psi' \in H^{\alpha/2}([0, T])$ , can be stated in the form

$$(2.24) \quad \psi \in Q^{\alpha/2}$$

where  $Q$  has been defined in (2.6). The change of variables  $y := x + 2bt^{1/2}$  is of no help for the proof of Theorem 2.2 since  $\mathbb{R}_+ \times [0, T]$  is transformed to the domain  $\{(y, t) : t \in [0, T], y > 2bt^{1/2}\}$ . However, we may assume  $a = 1$ , by a linear change of the  $t$ -variable.

Let us first obtain a representation for the solution of problem (2.20) (with  $a = 1, f = \phi = 0$ ) in terms of the fundamental solution  $K(x-y, t, s)$  for the equation

$$(2.25) \quad u_t - u_{xx} - t^{-1/2} b u_x = 0.$$

By taking Fourier-transforms we see that

$$(2.26) \quad \begin{aligned} K(x, t, s) &= \Gamma(x + 2b(t^{1/2} - s^{1/2}), t-s) \\ &= (4\pi)^{-1/2} (t-s)^{-1/2} \exp\left(-\frac{(x+2b(t^{1/2} - s^{1/2}))^2}{4(t-s)}\right). \end{aligned}$$

Using (2.1), (2.2), (2.14) and the fact that  $K$  satisfies equation (2.25) for  $x, t-s \neq 0$  one can easily check that  $K$  satisfies the same estimates as the fundamental solution of the heat equation

$$(2.27) \quad |D_x^j D_t^k K(x, t, s)| \leq c(t-s)^{-j/2-k} \exp(-c \frac{x^2}{t-s}).$$

Also note, that

$$(2.28) \quad \frac{\partial}{\partial s} K(x, t, s) + K_{xx}(x, t, s) + bs^{-1/2} K_x(x, t, s) = 0, \quad x, t-s \neq 0,$$

which follows from (2.26).

Proposition 2.1. Let  $\psi \in Q^{\alpha/2}$ . The solution of problem (2.20) (with  $f = \phi = 0$ ) can be represented in the form

$$(2.29) \quad u(x, t) = -2 \int_0^t K_x(x, t, s) \chi(s) ds$$

where  $\chi \in Q^{\alpha/2}$  is the solution of

$$(2.30) \quad \chi(t) = \psi(t) + 2 \int_0^t K_x(0, t, s) \chi(s) ds$$

and

$$(2.31) \quad |t^{1/2} \chi'|_{\alpha/2} \leq c |t^{1/2} \psi'|_{\alpha/2}.$$

Proof. We claim that the operator  $R$  defined by

$$(2.32) \quad (R\chi)(t) := -2 \int_0^t K_x(0, t, s) \chi(s) ds$$

is a strict contraction on the space  $Q$  with respect to the norm

$\|\chi\|_Q = |t^{1/2} \chi'|_{\alpha/2}$ . We have

$$-2 K_x(0, t, s) = \frac{b(t^{1/2} - s^{1/2})}{\pi^{1/2} (t-s)^{3/2}} \exp\left(-\frac{(b(t^{1/2} - s^{1/2}))^2}{t-s}\right)$$

and we rewrite  $R\chi$  in the form

$$\begin{aligned}
(R\chi)(t) &= \int_0^1 \frac{b}{\pi^{1/2}} \frac{1}{(1-s)^{1/2}(1+s)^{1/2}} \exp\left(-\frac{(b(1-s^{1/2}))^2}{1-s}\right) \chi(ts) ds \\
&=: \int_0^1 r(b,s) \chi(ts) ds.
\end{aligned}$$

Substituting  $z := (b(1-s^{1/2}))^2/(1-s)$  we see that

$$(2.33) \quad \|r(b, \cdot)\|_1 = \pi^{-1/2} \int_0^{b^2} \frac{b^2-z}{b^2+z} z^{-1/2} \exp(-z) dz < 1.$$

Since

$$t^{1/2}(R\chi)'(t) = \int_0^1 (r(b,s)s^{1/2})(ts)^{1/2} \chi'(ts) ds$$

this implies that  $R$  is a contraction on  $Q$ .

It remains to show that  $u$ , given by (2.29), satisfies the boundary condition  $u(0, \cdot) = \psi$ . We write (2.29) in the form

$$\begin{aligned}
u(x,t) &= \\
&\int_0^t \frac{x}{2\pi^{1/2}(t-s)^{3/2}} \exp\left(-\frac{(x+2b(t^{1/2}-s^{1/2}))^2}{4(t-s)}\right) \chi(t) ds \\
&+ \int_0^t \frac{x}{2\pi^{1/2}(t-s)^{3/2}} \exp\left(-\frac{(x+2b(t^{1/2}-s^{1/2}))^2}{4(t-s)}\right) (\chi(s) - \chi(t)) ds \\
&+ \int_0^t \frac{b(t^{1/2}-s^{1/2})}{\pi^{1/2}(t-s)^{3/2}} \exp\left(-\frac{(x+2b(t^{1/2}-s^{1/2}))^2}{4(t-s)}\right) \chi(s) ds \\
&=: \sum_{v=1}^3 I_v(x)
\end{aligned}$$

and obtain

$$\lim_{x \rightarrow 0} I_1(x) =$$

$$\chi(t) \lim_{x \rightarrow 0} \int_0^{t/x^2} \frac{1}{2\pi^{1/2}} s^{-3/2} \exp\left(-\frac{(1+2b((t/x^2)^{1/2} - (t/x^2-s)^{1/2}))^2}{4s}\right) ds =$$

$$\chi(t) \int_0^\infty \frac{1}{2\pi^{1/2}} s^{-3/2} \exp(-\frac{1}{4s}) ds = \chi(t) \quad .$$

$$|I_2(x)| \leq c \int_0^t x(t-s)^{-3/2} \exp(-c \frac{x^2}{t-s})(t-s)^{\alpha/2} |\chi|_{\alpha/2} ds \leq cx^\alpha |\chi|_{\alpha/2} \quad .$$

$$\lim_{x \rightarrow 0} I_3(x) = (R\chi)(t) = \psi(t) - \chi(t) \quad .$$

We now analyse the smoothness of the solution of problem (2.20) via the representation (2.29).

**Proposition 2.2.** Let  $u$  be defined by (2.29) with  $\chi \in \dot{H}^\gamma$ ,  $\gamma < 1/2$ , then

$$(2.34) \quad |u|_{\gamma,t} \leq c |\chi|_\gamma \quad .$$

**Proof.** Assume  $|\chi|_\gamma < 1$ . In particular, since  $\chi(0) = 0$ ,  $|\chi(t)| \leq t^\gamma$ . We let  $t-t' =: h \in (0, t/3)$  and write

$$\begin{aligned} & -\frac{1}{2} (u(x,t) - u(x,t')) = \\ & \int_{t-2h}^t K_x(x,t,s) (\chi(s) - \chi(t)) ds \\ & - \int_{t-2h}^{t'} K_x(x,t',s) (\chi(s) - \chi(t')) ds \\ & + \int_0^{t-2h} (K_x(x,t,s) - K_x(x,t',s)) (\chi(s) - \chi(t')) ds \\ & + \int_0^{2h} K_x(x,t,t-s) (\chi(t) - \chi(t')) ds \\ & + \int_0^{2h} (K_x(x,t,t-s) - K_x(x,t',t'-s)) \chi(t') ds \\ & + \int_0^{t'-2h} (K_x(x,t,t-2h-s) - K_x(x,t',t'-2h-s)) \chi(t') ds \\ & + \int_0^h K_x(x,t,h-s) \chi(t') ds \\ & =: \sum_{\nu=1}^7 I_\nu \quad . \end{aligned}$$

We set

$$(2.35) \quad \Lambda(t,s) := \frac{x+2b(t^{1/2} - s^{1/2})}{2(t-s)^{1/2}}$$

and with this abbreviation

$$\begin{aligned} K_x(x,t,s) &= - \frac{1}{2\pi^{1/2}} (t-s)^{-1} \Lambda(t,s) \exp(-\Lambda(t,s)^2) \\ &=: K_1(x,t,s) + K_2(x,t,s) \end{aligned}$$

where

$$\begin{aligned} K_1(x,t,s) &= - \frac{1}{4\pi^{1/2}} \frac{x}{(t-s)^{3/2}} \exp(-\Lambda(t,s)^2) \\ K_2(x,t,s) &= - \frac{1}{2\pi^{1/2}} \frac{b}{(t-s)^{1/2} (t^{1/2} + s^{1/2})} \exp(-\Lambda(t,s)^2) \end{aligned}$$

Note that, by (2.2),

$$(2.36) \quad \exp(-\Lambda(t,s)^2) \leq c \exp\left(-c \frac{x^2}{t-s}\right)$$

Using this,  $t-t' = h < t/3$  and (2.27) we estimate the integrals  $I_\nu$  as follows.

$$\begin{aligned} |I_1| &\leq c \int_{t-2h}^t (|K_1| + |K_2|) (t-s)^\gamma ds \\ &\leq c(h^\gamma + \int_{t-2h}^t t^{-1/2} (t-s)^{\gamma-1/2} ds) \leq ch^\gamma \end{aligned}$$

$|I_2|$  is estimated similarly.

$$\begin{aligned} |I_3| &\leq ch \int_0^{t-2h} |K_{xt}(x,\xi,s)| (t'-s)^\gamma ds \\ &\leq ch \int_0^{t-2h} (t'-s)^{-2+\gamma} dx \leq ch^\gamma \end{aligned}$$

$$\begin{aligned} |I_4| &\leq ch^\gamma \int_0^{2h} (|K_1| + |K_2|) \\ &\leq ch^\gamma (1 + \int_0^{2h} t^{-1/2} s^{-1/2} ds) \leq ch^\gamma \end{aligned}$$

Since  $|\frac{d}{d\lambda} (\lambda \exp(-\lambda^2))| < c$  and

$$(2.37) \quad \begin{aligned} & |\lambda(t, t-s) - \lambda(t', t'-s)| = \\ & cs^{-1/2} \left| \frac{s}{t^{1/2} + (t-s)^{1/2}} - \frac{s}{(t')^{1/2} + (t'-s)^{1/2}} \right| < \\ & cs^{1/2} t^{-1} h^{1/2} \end{aligned}$$

we obtain

$$\begin{aligned} |I_5| & < c(t')^\gamma \int_0^{2h} s^{-1} |\lambda(t, t-s) \exp(-\lambda(t, t-s)^2) - \\ & \quad \lambda(t', t'-s) \exp(-\lambda(t', t'-s)^2)| ds \\ & < ch^{1/2} t^{\gamma-1} \int_0^{2h} s^{-1/2} ds < ch^\gamma. \end{aligned}$$

Since

$$(2.38) \quad \begin{aligned} & |\lambda(t, t-2h-s) - \lambda(t', t'-2h-s)| = \\ & c(s+2h)^{-1/2} |t^{1/2} - (t-2h-s)^{1/2} - (t')^{1/2} + (t'-2h-s)^{1/2}| < \\ & c(s+2h)^{-1/2} h(t'-2h-s)^{-1/2} \end{aligned}$$

we obtain, for  $\gamma < 1/2$ ,

$$\begin{aligned} |I_6| & < c(t')^\gamma \int_0^{t'-2h} (s+2h)^{-1} |\lambda(t, t-2h-s) \exp(-\lambda(t, t-2h-s)^2) - \\ & \quad \lambda(t', t'-2h-s) \exp(-\lambda(t', t'-2h-s)^2)| ds \\ & < ct^\gamma \int_0^{t'-2h} (s+2h)^{-3/2} h(t'-2h-s)^{-1/2} ds \\ & < ct^\gamma h^{1/2} t^{-1/2} < ch^\gamma. \end{aligned}$$

$$|I_7| < c(t')^\gamma \int_0^h (t-h+s)^{-1} ds < ct^\gamma ht^{-1} < ch^\gamma.$$

**Proposition 2.3.** Let  $u$  be defined by (2.29) with  $x \in Q^\gamma$ ,  $\gamma < 1/2$ , then

$$(2.39) \quad |u_x|_{\gamma, t} < c |t^{1/2} x'|_\gamma.$$

Proof. Assume  $|t^{1/2}\chi'|_Y < 1$ . By (2.28) we have

$$\begin{aligned} u_x(x,t) &= -2 \int_0^t K_{xx}(x,t,s)\chi(s)ds = \\ &= -2 \int_0^t K(x,t,s)\chi'(s)ds + 2b \int_0^t K_x(x,t,s)s^{-1/2}\chi(s)ds \\ &=: v(x,t) + w(x,t) . \end{aligned}$$

Since, by Lemma 2.4,  $|t^{-1/2}\chi|_Y < c$ , Proposition 2.2 implies  $|w|_{Y,t} < c$ .

Let  $t-t' =: h \in (0, t/3)$ . As in the proof of Proposition 2.2 we write the difference  $-\frac{1}{2}(v(x,t) - v(x,t'))$  in the form  $\sum_{\nu=1}^7 J_\nu$  where the integrals  $J_\nu$  are defined as  $I_\nu$  but with  $K_x$  replaced by  $K$  and  $\chi$  by  $\chi'$ . Using  $t-t' = h < t/3$ , (2.27) and the inequalities (c.f. Lemma 2.3)

$$|\chi'(t)| < ct^{-1/2+\gamma}$$

$$|\chi'(t) - \chi'(s)| < c(t-s)^\gamma s^{-1/2}, \quad s < t ,$$

we estimate the integrals  $J_\nu$  as follows.

$$|J_1| < c \int_{t-2h}^t (t-s)^{-1/2} (t-s)^\gamma s^{-1/2} ds < ch^\gamma .$$

$|J_2|$  is estimated similarly.

$$\begin{aligned} |J_3| &< ch \int_0^{t-2h} |K_t(x, \xi, s)| (t'-s)^\gamma s^{-1/2} ds \\ &< ch \int_0^{t-2h} (t-s)^{-3/2+\gamma} s^{-1/2} ds < ch^\gamma . \end{aligned}$$

$$|J_4| < ch^\gamma (t')^{-1/2} \int_0^{2h} s^{-1/2} ds < ch^\gamma .$$

With  $\Lambda$  defined by (2.35) we have, using (2.37),

$$\begin{aligned} |J_5| &< c(t')^\gamma s^{-1/2} \int_0^{2h} s^{-1/2} |\exp(-\Lambda(t, t-s)^2) - \exp(-\Lambda(t', t'-s)^2)| ds \\ &< ct^{\gamma-1/2} \int_0^{2h} t^{-1} h^{1/2} ds < ch^\gamma . \end{aligned}$$

By (2.38),

$$\begin{aligned}
& |\exp(-A(t, t-2h-s)^2) - \exp(-A(t', t'-2h-s)^2)| \\
& \leq c(s+2h)^{-1/2} h(t'-2h-s)^{-1/2} |A(\xi, \xi-2h-s) \exp(-A(\xi, \xi-2h-s)^2)| \\
& \leq c(s+2h)^{-1/2} h(t'-2h-s)^{-1/2} \left| \frac{x}{(s+2h)^{1/2}} \exp\left(-c \frac{x^2}{s+2h}\right) + (s+2h)^{1/2} (t')^{-1/2} \right|.
\end{aligned}$$

It follows that

$$\begin{aligned}
|J_6| & \leq c(t')^{-1/2+\gamma} \int_0^{t'-2h} (s+2h)^{-1/2} |\exp(-A(t, t-2h-s)^2) - \\
& \quad \exp(-A(t', t'-2h-s)^2)| ds \\
& \leq ct^{-1/2+\gamma} h \left| \int_0^{t'-2h} \frac{x}{(s+2h)^{-3/2}} \exp\left(-c \frac{x^2}{s+2h}\right) (t'-2h-s)^{-1/2} ds \right. \\
& \quad \left. + t^{-1/2} \int_0^{t'-2h} (s+2h)^{-1/2} (t'-2h-s)^{-1/2} ds \right| \\
& \leq ct^{-1/2+\gamma} ht^{-1/2} \leq ch^\gamma. \\
|J_7| & \leq c(t')^{-1/2+\gamma} \int_0^h (t-h+s)^{-1/2} dx \\
& \leq ct^{-1/2+\gamma} ht^{-1/2} \leq ch^\gamma.
\end{aligned}$$

**Proposition 2.4.** For  $\psi \in Q^{\alpha/2}$ , the solution of problem (2.20) (with  $f = \phi = 0$ ) satisfies

$$(2.40) \quad |u_x|_{\alpha, x} \leq c |t^{1/2} \psi'|_{\alpha/2}.$$

**Proof.** Assume  $|t^{1/2} \psi'|_{\alpha/2} \leq 1$ . Since  $v := u_x$  satisfies (2.25) and  $v(\cdot, 0) = 0$ ,  $v(0, \cdot) = u_x(0, \cdot)$  we have

$$v(x, t) = -2 \int_0^t K_x(x, t, s) \bar{\chi}(s) ds$$

with  $\bar{\chi}$  the solution of

$$\bar{\chi}(t) = u_x(0, t) + 2 \int_0^t K_x(0, t, s) \bar{\chi}(s) ds.$$

By Proposition 2.3,  $|u_x(0, \cdot)|_{\alpha/2} \leq c$ , and the proof of Proposition 2.1, in particular (2.33), shows that this implies  $|\bar{\chi}|_{\alpha/2} \leq c$ .

Let  $x-x' =: h > 0$  and assume that  $t > h^2$ . We write

$$\begin{aligned}
 & -\frac{1}{2} (v(x,t) - v(x',t)) = \\
 & \int_{t-h^2}^t K_x(x,t,s) (\bar{\chi}(s) - \bar{\chi}(t)) ds \\
 & - \int_{t-h^2}^t K_x(x',t,s) (\bar{\chi}(s) - \bar{\chi}(t)) ds \\
 & + \int_0^{t-h^2} (K_x(x,t,s) - K_x(x',t,s)) (\bar{\chi}(s) - \bar{\chi}(t)) ds \\
 & + \int_0^t (K_x(x,t,s) - K_x(x',t,s)) \bar{\chi}(t) ds \\
 & =: \sum_{\nu=1}^4 I_\nu .
 \end{aligned}$$

Using  $h^2 < t$ , (2.27) and  $|\bar{\chi}(t)| \leq ct^{\alpha/2}$  we estimate these integrals as follows.

$$|I_1| \leq c \int_{t-h^2}^t (t-s)^{-1} (t-s)^{\alpha/2} ds \leq ch^\alpha .$$

$|I_2|$  is estimated similarly.

$$\begin{aligned}
 |I_3| & \leq ch \int_0^{t-h^2} |K_{xx}(\xi, t, s)| (t-s)^{\alpha/2} ds \\
 & \leq ch \int_0^{t-h^2} (t-s)^{-3/2+\alpha/2} ds \leq ch^\alpha .
 \end{aligned}$$

Set  $I(x) := \int_0^t K_x(x,t,s) ds$ . By (2.28) we have

$$\begin{aligned}
 |I'(x)| & = \left| \int_0^t K_{xx}(x,t,s) ds \right| \\
 & \leq |K(x,t,\cdot)|_0^t + c \int_0^t |K_x(x,t,s)| s^{-1/2} ds \\
 & \leq c(t^{-1/2} + \int_0^t \frac{x}{(t-s)^{3/2}} \exp(-c \frac{x^2}{t-s}) s^{-1/2} ds \\
 & \quad + \int_0^t \frac{1}{(t-s)^{1/2} (t^{1/2} + s^{1/2})} s^{-1/2} ds) \\
 & \leq ct^{-1/2} .
 \end{aligned}$$

It follows that

$$|I_4| \leq ct^{\alpha/2} ht^{-1/2} \leq ch^{\alpha}.$$

Combining the above estimates we have

$$|v(x,t) - v(x',t)| \leq c|x-x'|^{\alpha}, \quad |x-x'|^2 \leq t.$$

Since  $v(\cdot, 0) = 0$  and, by Proposition 2.3,  $|v|_{\alpha/2, t} \leq c$ , this inequality holds for  $|x-x'|^2 > t$  too.

Proof of Theorem 2.2. As we already remarked we may assume  $f = \phi = 0$ . For  $\psi \in Q^{\alpha/2}$  it follows from Propositions 2.1 - 2.4 that

$$(2.41) \quad \|u, u_x\|_{\alpha, \alpha/2} \leq c|t^{1/2}\psi'|_{\alpha/2}.$$

To complete the proof of assertions (2.22), (2.23) we write the solution of problem (2.20) in the form  $u = v + w$  where  $v$  and  $w$  are solutions of the problems

$$(2.42) \quad \begin{cases} v_t - v_{xx} = t^{-1/2} \overline{bu}_x, & (x,t) \in \mathbb{R} \times [0, T], \\ v(\cdot, 0) = 0 \end{cases}$$

$$(2.43) \quad \begin{cases} w_t - w_{xx} = 0, & (x,t) \in \mathbb{R}_+ \times [0, T], \\ w(\cdot, 0) = 0 \\ w(0, \cdot) = \psi - v(0, \cdot). \end{cases}$$

Here,  $\overline{u}_x$  denotes a smooth extension of  $u_x$  to the domain  $\mathbb{R} \times [0, T]$ . By (2.41) and Theorem 2.1,

$$\|v, v_x, t^{1/2}v_{xx}, w, w_x\|_{\alpha, \alpha/2} \leq c|t^{1/2}\psi'|_{\alpha/2}.$$

Also note that, by (2.10),

$$\lim_{t \rightarrow 0} t^{1/2}(\psi'(t) - v_t(0, t)) = (\lim_{t \rightarrow 0} t^{1/2}\psi'(t)) - b\phi_x(0) = 0,$$

and it follows from  $|t^{1/2}v_t|_{\alpha, \alpha/2} \leq |t^{1/2}v_{xx}|_{\alpha, \alpha/2} + |\overline{bu}_x|_{\alpha, \alpha/2}$  that  $\tilde{\psi} := \psi - v(0, \cdot) \in Q^{\alpha/2}$ . Therefore it remains to show that

$$(2.44) \quad |t^{1/2}w_{xx}|_{\alpha, \alpha/2} \leq c|t^{1/2}\tilde{\psi}'|_{\alpha/2}$$

$$(2.45) \quad \lim_{t \rightarrow 0} t^{1/2} w_{xx}(x, t) = 0, \quad x \in R_+.$$

Assume  $|t^{1/2} \tilde{\psi}'|_{\alpha/2} < 1$ . We have

$$(2.46) \quad \begin{aligned} |w_{xx}(x, t)| &= |w_t(x, t)| = |2 \int_0^t \Gamma_x(x, t-s) \tilde{\psi}'(s) ds| \\ &< c \int_0^t \frac{x}{(t-s)^{3/2}} \exp(-c \frac{x^2}{t-s}) s^{-1/2+\alpha/2} ds < c t^{-1/2+\alpha/2} \end{aligned}$$

which proves (2.45).

Let  $x-x' =: h > 0$  and assume that  $h^2 < t$ ,  $h < x/2$ . We have

$$\begin{aligned} |w_{xx}(x, t) - w_{xx}(x', t)| &= \\ |2 \int_0^t (\Gamma_x(x, t-s) - \Gamma_x(x', t-s)) \tilde{\psi}'(s) ds| &< \\ ch \int_0^t |\Gamma_{xx}(\xi, t-s)| s^{-1/2+\alpha/2} ds &< \\ ch \int_0^t (t-s)^{-3/2} \exp(-c \frac{x^2}{t-s}) s^{-1/2+\alpha/2} ds &< \\ ct^{-1/2} h^\alpha. \end{aligned}$$

In combination with (2.46) this shows that

$$|t^{1/2} w_{xx}|_{\alpha, x}, \quad |t^{1/2-\alpha/2} w_{xx}|_\infty < c.$$

To complete the proof of (2.44) note that

$$\begin{aligned} |w_{xx}|_{\alpha/2, t, R_+ \times [t, T]} &< \\ c(|w_{xx}(\cdot, t)|_\alpha + |\tilde{\psi}'|_{\alpha/2, [t, T]}) &< ct^{-1/2} \end{aligned}$$

and apply Lemma 2.3.

### 3. Extension to variable coefficients

#### 3.1. Reduction of the problem

We show in this section that it suffices to prove Theorem 1.1 for  $f \in H^{0,\alpha,\alpha/2}$  and  $\phi = \psi_v = 0$ .

Let  $\bar{b}, \bar{f}, \bar{\phi}$  be smooth extensions of the functions  $b, f, \phi$  to the domains  $R \times [0, T]$  and  $R$  respectively and set  $\tilde{\phi}(x, t) := \bar{\phi}(x)$ . Let  $v$  be the solution of the problem

$$(3.1) \quad \begin{cases} v_t - v_{xx} = t^{-1/2}(\bar{b} \tilde{\phi}_x + \bar{f}), & (x, t) \in R \times [0, T] \\ v(\cdot, 0) = \bar{\phi} \end{cases}$$

and set

$$V(x, t) := x(\psi_1(t) - v(1, t)) + (1-x)(\psi_0(t) - v(0, t)) .$$

Then the solution of problem (1.1) can be written in the form

$$(3.2) \quad u = v + V + w$$

where  $w$  is the solution of the problem

$$(3.3) \quad \begin{cases} w_t - aw_{xx} - t^{-1/2}bw_x = t^{-1/2}g, & (x, t) \in \Omega_T := [0, 1] \times [0, T] \\ w(x, 0) = 0, & x \in [0, 1] \\ w(v, t) = 0, & t \in [0, T], v = 0, 1 \end{cases}$$

with

$$g := t^{1/2}(a-1)v_{xx} + b(v_x - \tilde{\phi}_x) - t^{1/2}v_t + bv_x .$$

By Theorem 2.1 and the definition of  $V$  we have

$$(3.4) \quad \|v, v_x, t^{1/2}v_{xx}, t^{1/2}v_t, v, v_x, t^{1/2}v_t\|_{\alpha, \alpha/2} \leq c(\|f\|_{\alpha, \alpha/2} + \|\phi, \phi'\|_{\alpha} + \sum_{v=0}^1 \|\psi_v\|_{\alpha/2}) ,$$

where  $c$  depends on  $\|b\|_{\alpha, \alpha/2}$ . Moreover,

$$(3.5) \quad \lim_{t \rightarrow 0} t^{1/2} v_{xx}(x, t) = 0, \quad x \in [0, 1].$$

This, together with the compatibility conditions (1.5) and the equation (3.1), implies

$$\begin{aligned} \lim_{t \rightarrow 0} g(x, t) &= \lim_{t \rightarrow 0} (-t^{1/2} v_t(x, t) + b(x, t) v_x(x, t)) = \\ &= \lim_{t \rightarrow 0} (-t^{1/2} (x \psi_1'(t) + (1-x) \psi_0'(t))) + \\ &+ x(\bar{b}(1, 0) \phi'(1) + \bar{f}(1, 0)) + (1-x)(\bar{b}(0, 0) \phi'(0) + \bar{f}(0, 0)) \\ &+ b(x, 0)((\psi_1(0) - v(1, 0)) - (\psi_0(0) - v(0, 0))) = 0. \end{aligned}$$

By (3.4) and the assumptions (1.3) and (1.4) on  $a, b, \phi, \psi_v$  it follows that  $g \in H^{\alpha, \alpha/2}$ .

In view of the above reductions, Theorem 1.1 is a consequence of  
Theorem 3.1. For  $g \in H^{\alpha, \alpha/2}(\Omega_T)$ , the solution of problem (3.3) satisfies

$$(3.6) \quad \|w, w_x t^{1/2} w_{xx}\|_{\alpha, \alpha/2, \Omega_T} \leq c \|g\|_{\alpha, \alpha/2, \Omega_T}$$

where  $c$  depends on  $\alpha, a_0, \|a, b\|_{\alpha, \alpha/2, \Omega_T}$  and  $T$ . Moreover we have

$$(3.7) \quad \lim_{t \rightarrow 0} t^{1/2} w_{xx}(x, t) = 0.$$

### 3.2. Auxiliary lemmas

For  $\lambda := 1/J$ ,  $J \in \mathbb{N}$ , we set  $x_j := j\lambda$ ,  $j = 0, \dots, J$ , and choose cut off functions  $\eta_j$  with the following properties

$$(3.8) \quad \left\{ \begin{array}{l} 0 < \eta_j(x) < 1 \\ \sum_j \eta_j(x)^2 = 1, \quad x \in [0,1] \\ x_j \in \text{supp } \eta_j \\ |\text{supp } \eta_j| < c\lambda \\ \|D^k \eta_j\|_\infty < c\lambda^{-k} \end{array} \right.$$

Note that, by the third and fourth property, the supports of at most  $c$  (independent of  $\lambda$ )  $\eta_j$ 's can have a common intersection. We set

$$M_j(x, t) := \eta_j(x) \quad .$$

and for simplicity of notation

$$\|\cdot\|_T := \|\cdot\|_{\alpha, \alpha/2, \Omega_T} \quad .$$

$$(3.9) \quad \begin{array}{l} \text{Lemma 3.1. For } f = \sum_j f_j \text{ with } \text{supp } f_j \subseteq \text{supp } M_j, \\ \|f\|_T < c \sup_j \|f_j\|_T \end{array}$$

Proof. Let us prove the estimate, e.g., for  $\|\cdot\|_{\alpha, x}$ . For  $x, x'$  set  $I := \{j : x \in \text{supp } \eta_j \text{ or } x' \in \text{supp } \eta_j\}$ . Since by (3.8)  $|I| < c$ , we have

$$\begin{aligned} |f(x, t) - f(x', t)| &= \left| \sum_{j \in I} f_j(x, t) - f_j(x', t) \right| \\ &< c|x-x'|^\alpha \sup_{j \in I} \|f_j\|_{\alpha, x} \quad . \end{aligned}$$

Lemma 3.2. For  $f \in H^{\alpha, \alpha/2}(\Omega_T)$  with  $T < \lambda^2$ ,

$$(3.10) \quad \sup_j \|M_j f\|_T < c \|f\|_T \quad ,$$

where the constant  $c$  does not depend on  $\lambda$  and  $T$ .

This follows from (3.8) and the inequality

$$(3.11) \quad \|uv\|_Y \leq \|u\|_\infty \|v\|_Y + \|u\|_Y \|v\|_\infty.$$

**Lemma 3.3.** If  $w(\cdot, 0) = w_x(\cdot, 0) = 0$  and  $\|w\|_{1/2+\alpha/2, t, \Omega_T} \leq 1$ , then

$$(3.12) \quad \|(D_x^k M_j) t^{l/2} D_x^m w\|_T \leq c \lambda^{-k} T^{(l-m+1)/2}, \quad m = 0, 1,$$

where the constant  $c$  does not depend on  $\lambda$  and  $T$ .

**Proof.** Let us prove, e.g., the inequality

$$\|w\|_{\alpha, x, \Omega_T} \leq c T^{1/2}.$$

For  $|x-x'| \leq T^{1/2}$  we have

$$\begin{aligned} |w(x, t) - w(x', t)| &\leq |x-x'| \|w_x\|_\infty \\ &\leq |x-x'| T^{1/2-\alpha/2} T^{\alpha/2} \|w_x\|_{\alpha/2, t, \Omega_T} \end{aligned}$$

and for  $|x-x'| > T^{1/2}$ ,

$$\begin{aligned} |w(x, t) - w(x', t)| &\leq 2|x-x'| T^{-\alpha/2} \|w\|_\infty \\ &\leq 2|x-x'| T^{-\alpha/2} T^{1/2+\alpha/2} \|w\|_{1/2+\alpha/2, t, \Omega_T}. \end{aligned}$$

All other inequalities can be easily deduced, using (3.11), (3.8) and the inequality

$$(3.13) \quad \|x\|_{Y', [0, T]} \leq T^{\gamma-\gamma'} \|x\|_{Y, [0, T]}$$

for  $x \in H^{\circ\gamma}([0, T])$  and  $\gamma' < \gamma$ .

### 3.3. Proof of the main theorem

Denote by  $L$  the differential operator

$$L : w \mapsto w_t - a w_{xx} - t^{-1/2} b w_x.$$

To prove Theorem 3.1 we define an approximate right inverse  $\tilde{L}^{-1}$  to  $L$  by

$$(3.14) \quad \tilde{L}^{-1}(t^{-1/2} g) = \sum_j M_j L_j^{-1}(M_j t^{-1/2} g)$$

where  $w_j := L_j^{-1}(M_j t^{-1/2} g)$  is the solution of the boundary value problem

$$(3.15) \quad \begin{cases} (w_j)_t - a(x_j, 0)(w_j)_{xx} - t^{-1/2} b(x_j, 0)(w_j)_x = M_j t^{-1/2} g & \text{on } \Omega_j, \\ w_j = 0 & \text{on } \partial\Omega_j \end{cases}$$

with  $\Omega_j := \mathbb{R}_+ \times [0, T]$  if  $0 \in \text{supp } \eta_j$  and  $\Omega_j := (-\infty, 1) \times [0, T]$  otherwise.

Proposition 3.1. Set  $\tilde{w} := \tilde{L}^{-1}(t^{-1/2} g)$ . Then, for  $g \in \dot{H}^{\alpha, \alpha/2}(\Omega_T)$  with  $T < \lambda^2$ , we have  $\tilde{w}, \tilde{w}_x, t^{1/2} \tilde{w}_{xx} \in \dot{H}^{\alpha, \alpha/2}(\Omega_T)$  and

$$(3.16) \quad \|\tilde{w}, \tilde{w}_x, t^{1/2} \tilde{w}_{xx}\|_T \leq c \|g\|_T$$

where the constant  $c$  does not depend on  $\lambda$  and  $T$ .

Proof. Assume that  $T < \lambda^2$  and  $\|g\|_T < 1$ . By Lemma 3.2,  $\|M_j g\|_T \leq c$ .

Applying Theorem 2.2 to problem (3.1), it follows that  $w_j, (w_j)_x,$

$(t^{1/2} w_j)_{xx} \in \dot{H}^{\alpha, \alpha/2}(\Omega_T)$  and

$$(3.17) \quad \|w_j, (w_j)_x, (t^{1/2} w_j)_{xx}\|_T \leq c.$$

By Lemma 2.4 and (3.15) this implies in particular

$$(3.18) \quad \|w_j\|_{1/2+\alpha/2, t, \Omega_T} \leq \|t^{1/2} (w_j)_t\|_T \leq c.$$

Since  $\tilde{w} = \sum_j M_j w_j$  we obtain, using Lemmas 3.1 - 3.3,

$$\|\tilde{w}\|_T \leq c \sup_j \|M_j w_j\|_T \leq c T^{1/2}.$$

$$\|\tilde{w}_x\|_T \leq c \sup_j (\|(M_j)_x w_j\|_T + \|M_j (w_j)_x\|_T) \leq c.$$

$$\|(t^{1/2} \tilde{w})_{xx}\|_T \leq$$

$$c \sup_j (\|(M_j)_{xx} t^{1/2} w_j\|_T + \|(M_j)_x (t^{1/2} w_j)_x\|_T + \|(t^{1/2} w_j)_{xx}\|_T) \leq c.$$

**Proposition 3.2.** The operator

$$S : g \mapsto t^{1/2} L \tilde{L}^{-1} (t^{-1/2} g)$$

maps  $H^{\alpha, \alpha/2}(\Omega_T)$  into itself. There exist  $\lambda_0$  and  $T_0 < \lambda_0^2$  such that

$$(3.19) \quad \|Sg - g\|_{T_0} < \frac{1}{2} \|g\|_{T_0}.$$

**Proof.** The first assertion of the Proposition is clear from Proposition 3.1.

We write

$$\begin{aligned} & t^{1/2} L \tilde{L}^{-1} (t^{-1/2} g) - g = \\ & t^{1/2} \left( \sum_j (L(M_j w_j) - M_j L w_j) + \sum_j M_j (L - L_j) w_j \right) = \\ & t^{1/2} \sum_j a(M_j)_{xx} w_j + 2 t^{1/2} \sum_j a(M_j)_x (w_j)_x + \sum_j b(M_j)_x w_j + \\ & t^{1/2} \sum_j M_j (a - a(x_j, 0)) (w_j)_{xx} + \sum_j M_j (b - b(x_j, 0)) (w_j)_x \\ & =: \sum_{\nu=1}^5 I_\nu, \end{aligned}$$

where we used that

$$t^{1/2} \sum_j M_j L_j w_j = t^{1/2} \sum_j M_j L_j L_j^{-1} (M_j t^{-1/2} g) = g$$

since  $\sum_j (M_j)^2 = 1$ . Assuming  $T < \lambda^2$ ,  $\|g\|_T < 1$  and using (3.17), (3.18) and Lemmas 3.1 - 3.3 we estimate these terms as follows.

$$\|I_1\|_T < c \sup_j \|a\|_T \| (M_j)_{xx} t^{1/2} w_j \|_T < c \lambda^{-2} T.$$

$$\|I_2\|_T < c \sup_j \|a\|_T \| (M_j)_x (t^{1/2} w_j)_x \|_T < c \lambda^{-1} T^{1/2}.$$

$$\|I_3\|_T < c \sup_j \|b\|_T \| (M_j)_x w_j \|_T < c \lambda^{-1} T^{1/2}.$$

$$\begin{aligned}
& \|I_4\|_T < \\
& c \sup_j (\|a - a(x_j, 0)\|_{\infty, \text{supp } M_j} \|M_j(t^{1/2} w_j)_{xx}\|_T + \\
& \|a\|_T \|M_j(t^{1/2} w_j)_{xx}\|_{\infty}) < \\
& c (\lambda^\alpha + T^{\alpha/2}) .
\end{aligned}$$

$$\begin{aligned}
& \|I_5\|_T < \\
& c \sup_j (\|b - b(x_j, 0)\|_{\infty, \text{supp } M_j} \|M_j(w_j)_x\|_T + \|b\|_T \|M_j(w_j)_x\|_{\infty}) < \\
& c (\lambda^\alpha + T^{\alpha/2}) .
\end{aligned}$$

Combining these estimates we have

$$\| \sum_{v=1}^5 I_v \|_T < c(\lambda^{-2} T + \lambda^{-1} T^{1/2} + \lambda^\alpha + T^{\alpha/2})$$

which proves the Proposition with  $\lambda_0 := (8c)^{-1/\alpha}$ ,  $T_0 := (\lambda_0/(8c))^2$ .

By Proposition 3.2, the operator  $S$  is boundedly invertible on  $H^{\alpha, \alpha/2}(\Omega_{T_0})$ . It follows that, for  $g \in H^{\alpha, \alpha/2}(\Omega_{T_0})$ ,  $\tilde{L}^{-1}(t^{-1/2} S^{-1} g)$  is a solution of problem (3.3). Therefore, Proposition 3.1 proves Theorem 3.1 with  $T = T_0$ . Using the standard results for linear parabolic equations with smooth coefficients [L, Thm. 5.2, p. 320] the estimate (3.6) remains valid for any finite rectangle  $\Omega_T$ . In view of the reductions made in section 3.1, the proof of Theorem 1.1 is complete.

#### Acknowledgement

We thank Professor J. A. Nohel for many fruitful discussions, in particular concerning material in section 2.3.

#### REFERENCE

- [L] Ladyženskaja, O. A., Solonnikov, V. A., Uralčeva, N. N., Linear and quasilinear equations of parabolic type, Translations of Mathematical monographs, vol. 23, 1968.

KH/jvs

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER #2453	2. GOVT ACCESSION NO. AD-A125 243	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Linear Parabolic Equations with a Singular Lower Order Coefficient		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Klaus Höllig		8. CONTRACT OR GRANT NUMBER(s) MCS-7927062, Mod. 2. DAAG29-80-C-0041
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 1 - Applied Analysis
11. CONTROLLING OFFICE NAME AND ADDRESS  See Item 18 below		12. REPORT DATE November 1982
		13. NUMBER OF PAGES 29
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)  UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES U. S. Army Research Office P. O. Box 12211 Research Triangle Park North Carolina 27709  National Science Foundation Washington, DC 20550		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  parabolic equations, linear, singular, regularity		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Let $a, b$ belong to the Hölder class $H^{\alpha, \alpha/2}([0,1] \times [0,T])$ with $\alpha \in (0,1)$ and $a \geq a_0 > 0$ . It is shown that for the solution of the problem $u_t - au_{xx} - t^{-1/2}bu_x = t^{-1/2}f, \quad (x,t) \in [0,1] \times [0,T],$ $u(\cdot, 0) = \phi$ $u(v, \cdot) = \psi_v, \quad v = 0, 1,$ the estimate		

ABSTRACT (continued)

$$\|u, u_x, t^{1/2} u_{xx}\|_{\alpha, \alpha/2} \leq$$

$$c\{\|f\|_{\alpha, \alpha/2} + \|\phi, \phi'\|_{\alpha} + \sum_{v=0}^1 \|\psi_v, t^{1/2} \psi'_v\|_{\alpha/2}\}$$

holds if the data  $f, \phi, \psi$  satisfy the appropriate compatibility conditions. Here  $\|\cdot\|_{\alpha, \alpha/2}, \|\cdot\|_{\alpha}, \|\cdot\|_{\alpha/2}$  denote the norms of the Hölder classes  $H^{\alpha, \alpha/2}([0,1] \times [0,T]), H^{\alpha}([0,1])$  and  $H^{\alpha/2}([0,T])$  respectively and the constant  $c$  depends on  $\alpha, a_0, \|a, b\|_{\alpha, \alpha/2}$  and  $T$ . The result extends to quasilinear problems with  $a, b$  and  $f$  depending on  $x, t$  and  $u$ .